

Intermountain End-Stage Renal Disease Network, Inc.
(Network #15)
2008 CPM Results and Network Goals

CPM Element	Network #15 Results	U.S. Results
<p>QI 5. Dialysis Adequacy</p> <p><i>CMS Goal:</i> Attain/maintain 80% of patients with URR \geq 65%</p> <p>Attain/maintain 80% of patients with Kt/V \geq 1.2</p> <p>Our Network stretch goal is 95% for both indicators</p>	<p>2007 CPM:</p> <p>89% of sampled pts. in NW 15 had a URR \geq 65%</p> <p>91% of sampled pts in NW15 had a Kt/V of \geq 1.2</p>	<p>U.S. 2007 CPM:</p> <p>89%</p> <p>91%</p>

CPM Element	Network #15 Results	U.S. Results
<p>QI 6. Anemia Management</p> <p><i>CMS Goal:</i> Attain/maintain 70% of patients with mean hemoglobin \geq 11gm/dL</p> <p>Our Network stretch goal is 85%</p>	<p>2007 CPM:</p> <p>83% of sampled pts in NW 15 were reported to have a mean Hgb of \geq 11.0 gm/dL</p>	<p>U.S. 2007 CPM:</p> <p>82%</p>

CPM Element	Network #15 Results	U.S. Results
<p>QI 10. Nutrition</p> <p><i>CMS Goal:</i> 35% of patients to have a mean serum albumin \geq 4.0/3.7 gm/dL (BCG/BCP).</p> <p>Our Network stretch goal is 100%!</p>	<p>2007 CPM:</p> <p>36% of pts sampled in NW 15 were reported to have a serum albumin of \geq 4.0 gm/dL</p>	<p>U.S. 2007 CPM:</p> <p>34%</p>

CPM Element	Network #15 Results	U.S. Results
<p>QI 8. Vascular Access</p> <p><i>CMS Goal:</i> Increase the use of fistulas: At least 45 % of <i>prevalent</i> hemodialysis patients over 18 years of age should have a native AV fistula.</p> <p>Our Network stretch goal is 65%</p> <p>At least 50% of <i>incident</i> patients over 18 years of age should have a native AV fistula.</p> <p>Our Network stretch goal is 75%</p>	<p>2007 CPM:</p> <p>57% of sampled <i>prevalent</i> pts in NW 15 were reported to have a native AVF</p> <p>45% of sampled <i>incident</i> patients were reported to have a native AVF</p>	<p>U.S. 2007 CPM:</p> <p>49%</p> <p>41%</p>
<p>QI 9.</p> <p><i>CMS Goal:</i> Monitoring AV grafts for stenosis: should occur for 100% of all hemodialysis patients.</p>	<p>62% of pts sampled in NW 15 with an AVG were monitored for Stenosis</p>	<p>71%</p>
<p>QI 7.</p> <p><i>CMS Goal:</i> Decrease catheter use</p> <p>Less than 10% of prevalent hemodialysis patients over 18 years of age should be maintained on catheters for 90 days or longer.</p>	<p>23% of sampled pts in NW 15 were reported to have a catheter access \geq 90 days</p>	<p>21%</p>
<p>Ranges for other lab values that are audited by the State Survey Agencies (SSAs) are indicated on the Measures Assessment Tool included with the Conditions for Coverage. You may link to the Conditions document from the Network #15 web site @: http://www.esrdnet15.org/</p>		